

QFT™ improves efficiency of TB contact investigations – allowing you to focus your efforts

Imagine being able to focus only on individuals truly infected with TB...

Study Overview

In a high school TB contact investigation involving 349 contacts: ⁽¹⁾

- Out of 95 contacts who were TST (Tuberculin Skin Test) positive, only 4 were QFT™ positive.
- The 91 contacts who were TST positive—but QFT™ negative—were not given Isoniazid (INH) therapy, while the 4 contacts that were QFT positive were given INH therapy.
- After 3½ years of follow-up none of the 91 QFT™ negative—but TST positive—contacts had developed active TB.

...now it's possible with QFT™

Efficiency gains with QFT™

QFT™ Advantages	Limitations of TST (Tuberculin Skin Test) Programs
Need for one visit saves time and can improve adherence.	Problems with testing logistics commonly affect adherence to TST programs. <ul style="list-style-type: none"> • Requires two visits.
Unprecedented accuracy in detecting TB infection. <ul style="list-style-type: none"> • Unaffected by BCG vaccination.⁽⁴⁾ • Unaffected by most environmental non-tuberculous mycobacterium (NTM).⁽⁶⁾ 	TST accuracy adversely affected by previous BCG vaccination and non-tuberculous mycobacterial infections (NTM). <ul style="list-style-type: none"> • BCG can cause false positive TST responses in up to 20-40% of cases.⁽²⁾ • Up to 50% of TST responses can be due to NTM infections.⁽³⁾
Increased test accuracy provides confidence in initiating therapy and encourages therapy compliance.	Low compliance rates amongst TB contacts indicated for INH therapy. <ul style="list-style-type: none"> • Studies have shown that ~43% indicated TB contacts completed INH therapy.⁽⁶⁾

Numerous studies have shown that QFT™ has better correlation with the degree of exposure of contacts to their source case than the TST. ^(7,8,9,10,11)

In a large contact investigation involving >20,000 contacts exposed to a smear-positive supermarket employee:

“...TST results were associated with age while positive Interferon-γ responses were significantly associated with cumulative shopping time, most markedly for QFT-GIT. [the authors concluded]...Blood test results were associated with exposure, while the TST was not.” ⁽¹¹⁾

“QuantIFERON-TB Gold will eliminate the waste of precious health resources from inaccurate skin tests and improve public safety by preventing unnecessary treatment.”

Dr. Masae Kawamura, TB Controller for the City of San Francisco

Ordering Information

Catalogue Number	Product Description
0590 0301	QuantIFERON®-TB Gold In-Tube (Nil, TB Antigen, Mitogen tubes) 100 each
0597 0201	QuantIFERON®-TB Gold In-Tube Single Patient Sampling Pack (10 ct)
0594 0201	QuantIFERON®-TB Gold ELISA only

References

- Higuchi K, Harada N, Mori T, Sekiya Y. Use of QuantIFERON-TB Gold to investigate tuberculosis contacts in a high school. *Respirology*. 2007; 12(1):88-92.
- Farhat M, Greenaway C, Pai M, Menzies D. False-positive tuberculin skin tests: what is the absolute effect of BCG and non-tuberculous mycobacteria? *Int J Tuberc Lung Dis*. 2006; 10: 1192-204.
- von Reyn CF, Horsburgh CR, Olivier KN, et al. Skin test reactions to Mycobacterium tuberculosis purified protein derivative and Mycobacterium avium sensitin among health care workers and medical students in the United States. *Int J Tuberc Lung Dis*. 2001; 5 (12): 1122-1128.
- Mori T, Sakatani M, Yamagishi F, et al. Specific detection of tuberculosis infection: an interferon-gamma-based assay using new antigens. *Am J Respir Crit Care Med*. 2004; 170(1): 59-64.
- Anderson P, Munk ME, Pollock JM, Doherty TM. Specific immune-based diagnosis of TB. *Lancet*. 2000; 356: 1099-1104.
- Sprinson JE, Flood J, Fan CS, Shaw TA, Pascopella L, Young JA, Royce SE. Evaluation of tuberculosis contact investigations in California. *Int J Tuberc Lung Dis*. 2003; 7(12 Suppl 3):S363-8.
- Brock I, Munk ME, Kok-Jensen A, Andersen P. Performance of whole blood IFN-gamma test for tuberculosis diagnosis based on PPD or the specific antigens ESAT-6 and CFP-10. *Int J Tuberc Lung Dis* 2001; 5: 462-67.
- Funayama K, Tsujimoto A, Mori M, Yamamoto H, Fujiwara K, Nishimura T, Hasegawa N, Horiguchi I, Mori T, Marui E. Usefulness of QuantIFERON® TB-2G in contact investigation of a tuberculosis outbreak in a university. *Kekkaku* 2005; 80: 527-34.
- Brock I, Weldingh K, Lillebaek T, Follmann F, Andersen P. Comparison of tuberculin skin test and new specific blood test in tuberculosis contacts. *Am J Respir Crit Care Med* 2004; 170: 65-69.
- Diel R, Ernst M, Döschner G, et al. Avoiding the effect of BCG vaccination in detecting Mycobacterium tuberculosis infection with a blood test. *Eur Respir J* 2006; 28: 6-23.
- Arend SM, Thijsen SF, Leyten EM, Bouwman JJ, Franken WP, Koster BF, Cobelens FG, van Houte AJ, Bossink AW. Comparison of two interferon-gamma assays and tuberculin skin test for tracing tuberculosis contacts. *Am J Respir Crit Care Med*. 2007; 175(6):618-27.

For more information, please contact the Cellestis office nearest you or visit www.cellestis.com.

North America / South America
 Cellestis Inc.
 Email: customer.service@cellestis.com
 Fax: +1 661 775 7479
 Tel: +1 661 775 7480 (outside USA)
 Toll free: 800 519 4627 (USA only)

Europe / Middle East / Africa
 Cellestis Europe
 Email: europe@cellestis.com
 Fax: +49 6151 428 59 110
 Tel: +49 6151 428 59 0

Asia / Oceania
 Cellestis International
 Email: quantiferon@cellestis.com
 Fax: +61 3 9571 3544
 Tel: +61 3 9571 3500